



## A Pearl Millet: Effect On Climate Change Environment

Dr. Kumar Amit

Ph.D. (Science). PG Dept. of Botany, Maharaja College. VKSU. Ara. Bihar.

Corresponding Author: Dr. Kumar Amit

DOI -10.5281/zenodo.10988294

### Abstract:

*In common usage, climate change describes global warming—the ongoing increase in global average temperature—and its effects on Earth's climate system. Climate change in a broader sense also includes previous long-term changes to Earth's climate. The current rise in global average temperature is more rapid than previous changes and is primarily caused by humans burning fossil fuels.[3][4] Fossil fuel use, deforestation, and some agricultural and industrial practices add to greenhouse gases, notably carbon dioxide and methane. Greenhouse gases absorb some of the heat that the Earth radiates after it warms from sunlight. Larger amounts of these gases trap more heat in Earth's lower atmosphere, causing global warming.*

*The C4 grass pearl millet is one of the most drought-tolerant cereals and is primarily grown in marginal areas where annual rainfall is low and intermittent. It was domesticated in sub-Saharan Africa, and several studies have found that it uses a combination of morphological and physiological traits to successfully resist drought. This review explores the short-term and long-term responses of pearl millet that enable it to either tolerate, avoid, escape, or recover from drought stress. The response to short-term drought reveals fine-tuning of osmotic adjustment, stomatal conductance, and ROS scavenging ability, along with ABA and ethylene transduction. Equally important are longer-term developmental plasticity in tillering, root development, leaf adaptations, and flowering time that can help avoid the worst water stress and recover some of the yield losses via asynchronous tiller production.*

### Introduction:

C4 millets, including foxtail millet (*Setaria italica*), barnyard millet (*Echinochloa frumentacea* and *E. esculenta*), proso or common millet (*Panicum milieu*), and pearl millet (*Cenchrus americanus*) are extremely drought tolerant, and are relied on by farmers in marginal growing environments in many regions of the world. Pearl millet is one of the most

tolerant, being domesticated from its wild progenitor, *Cenchrus americanus* ssp. *violaceum* in the Sahel region of West Africa (Oumar et al., 2008; Burgarella et al., 2018), where annual rainfall varies between 200 to 600 mm (Bidinger et al., 1987). In this region, growing areas are characterized by a long dry season and a highly variable and short rainy season, conditions where few other grain crops would

consistently yield, or even survive (Debieu et al., 2018). Several pearl millet accessions have been found from even drier regions, including areas in the Mauritania and Mali regions with average rainfall less than 50 mm (Harlan, 1975; Pucher et al., 2015). Because of its drought-hardiness, pearl millet is pre-adapted for the predicted increased desertification that will result from climate change (Council, 2010), and has behavioural responses to drought damage that preserve at least some yield for the farmer. Recent breeding efforts have been directed at increasing yield and decreasing susceptibility to diseases (Serba et al., 2017), Pearl millet has the potential to play an important role in feeding the expanding world population, predicted to reach 9.1 billion by 2050, where, to meet the food demand of this huge population, cereal production needs to increase from 2.1 billion to up to 3 billion tons (Alexandratos, 2009). Making better use of marginal lands for crop cultivation will be an important component in increasing food production, and pearl millet will be a useful crop in these areas, as well as offering many lessons in tolerating drought that may be able to be translated to other crops. Importantly, breeders need to take the drought responses of pearl millet in the landscape in which it is best adapted

into consideration in their breeding efforts. Studies of the effects of drought on pearl millet include genetic analyses of variation in flowering time (Yadav et al., 2004). This review analyses the characteristics of pearl millet which foster drought resistance, including both short- and longer-term responses to drought. The great variation in pearl millet accessions that is available, as well as the feasibility of introgressions desirable characteristics from wild species, results in a huge range of possibilities for engineering drought tolerance/resistance in pearl millet.

#### **Materials and Methods:**

Pearl millet responses to drought resistance in pearl millet, drought studies have been conducted in each of the three main phases: vegetative, panicle development, and grain filling (Shivhare et al., 2020a) and these studies have identified resistance mechanisms involving drought avoidance, tolerance, escape, and recovery. These mechanisms can be grouped into short and long-term responses, as we discuss below (Figure 1). In addition, pearl millet is a C4 grass using the NADP-dependent malic enzyme (NADP-ME) pathway and is thus already preadapted for efficient water use in high-light environments (Pardo and VanBuren, 2021).




FIGURE 1 Short-term and long-term responses of pearl millet to adapt drought stress. ROS, Reactive oxygen species; HO, Hydroxyl radical; H<sub>2</sub>O<sub>2</sub>, Hydrogen peroxide; APX, Ascorbate peroxidase; GPX, Glutathione peroxidase; SOD, Superoxide dismutase.

**Short-term responses** Stomatal conductance Stomata are vital to the plant's existence, as they provide passage for gas and water exchange to conduct photosynthesis and transpiration (Li et al., 2017). Generally, stomatal closure is highly influenced by environmental conditions and is the first step for avoiding water loss in water-stressed conditions, but plants do it at the expense of gaining carbon dioxide for carbon fixation. Stomatal conductance in pearl millet has been implicated in

**Boosting Sustainability:**

Through offering a reduced dependence on synthetic fertilizers and pesticides, millets cultivation may also help promote a shift towards sustainable agriculture, diversifying crop rotations and avoiding the promotion of mono-cropping

systems. "The high carbon content of the crop residues makes them particularly important for maintaining and increasing soil carbon levels, important for sustainable cropping systems, and, where applicable, for providing forage, at the same time, for livestock," noted Dr. Aburto.

**Food System Divide:**

Millets are believed to be among the earliest domesticated plants, which have long served as traditional staple crops for millions of farmers, particularly in India, China, and Nigeria.

Notwithstanding the wide range of benefits that millets provide, they have largely been missing from the global food security agenda. In fact, in recent years, their production has gradually declined.

Experts point towards market distortions, a lack of appreciation of the

benefits of millets and policies that have favored the production of the so-called Big Three cereals - rice, wheat and maize, resulting in a "Food System Divide".

Joanna Kane-Potaka of ICRISAT, gave the example of India where "during the green revolution, high yielding varieties of rice and wheat were introduced and supported to scale out on a massive scale, to improve food security, while arguably, inadequate attention was paid to nutrition or environmental factors."

The problem is further compounded by changing dietary habits, high transaction costs and the challenges involved in accessing better markets; especially true for Africa.

"Farmers have therefore shifted to more remunerative crops grown to sell for profit and moved away from subsistence agriculture responding to changing consumer preferences and markets inputs," said Dr. Aburto.

### **2023: The Year of Millets:**

In declaring 2023 the International Year of Millets, the resolution calls on all stakeholders to provide support to "activities aimed at raising awareness of and directing policy attention to the nutritional and health benefits of millet consumption, and their suitability for cultivation under adverse and changing climatic conditions, while also directing

policy attention to improving value chain efficiencies."

Building on the experiences gained from past initiatives such as the 2016 International Year of Pulses and the 2021 International Year of Fruits and Vegetables, the UN agriculture agency is working to develop an action plan in partnership with external stakeholders, including farmers and research institutions.

"Actions taken will be aligned and supported via existing initiatives, such as the UN decade of action on Nutrition, 2016-2025 that provides an umbrella for a wide group of actors to work together to address malnutrition and other pressing nutrition issues," Dr. Aburto added.

In line with FAO's vision of a sustainable and food secure world for all, producing more and nutritious food for a growing population without overburdening land resources is a massive global challenge.

In the search for climate resilient solutions, millets could be the crucial link in the sustainable food supply chain.

Osmotic adjustment Osmotic adjustment is a major drought-tolerant mechanism in plants to maintain cell turgor, relative water content, cell expansion, photosynthesis, and continued stomatal conductance (Jones and Turner, 1980; Henson et al., 1982). Plants accumulate organic and inorganic

cellular components as osmolytes for lowering osmotic potential. In field conditions, pearl millet has been reported to adjust its osmotic potential successfully in response to water stress (Henson et al., 1982), like other drought-tolerant C3 and C4 grasses, such as upland rice (Lum et al., 2014), wheat genotypes (Hong-Bo et al., 2006), and sorghum (Blum and Ebercon, 1976). The biochemical basis of osmotic adjustment has both inorganic as well as organic components,

**Screening Techniques:**


- Spray uredospores collected from infector rows, on 25-30 days old crop, twice, at 25 and 35 days after sowing.
- Spreading of uredinia-bearing leaves among test plants 25-30 days old.

**Management:**

- Use of resistant hybrids/verieties.
- Sow the crop with the onset of monsoon.

- Destruction of collateral hosts like *Ischaemum pretosum* and *Panicum maximum* on the field bunds.
- Dusting of fine sulphur @ 17kg/ and two sprays of 0.2% Mancozeb at 15 days intervals

Osmotic adjustment is a major drought-tolerant mechanism in plants to maintain cell turgor, relative water content, cell expansion, photosynthesis, and continued stomatal conductance (Jones and Turner, 1980; Henson et al., 1982). Plants accumulate organic and inorganic cellular components as osmolytes for lowering osmotic potential. In field conditions, pearl millet has been reported to adjust its osmotic potential successfully in response to water stress (Henson et al., 1982), like other drought-tolerant C3 and C4 grasses, such as upland rice (Lum et al., 2014), wheat genotypes (Hong-Bo et al., 2006), and sorghum (Blum and Ebercon, 1976). The biochemical basis of osmotic adjustment has both inorganic as well as organic components.



**Fig:** (a) Frequency distribution of the 16,472 polymorphic SNPs across seven chromosomes; (b) Frequency distribution of the minor alleles for each SNP markers scored in a population sample of 109 genotypes; (c) Transition and transversion mutations of GBS-SNPs detected among 109 pearl millet hybrid parental lines; (d) Identity-by-state (IBS) based genetic distance between pairs of hybrid parental lines.

Smut is widely distributed in all the pearl millet growing areas in the world. Systemic symptoms as chlorosis generally appear on the second leaf and all the subsequent leaves and panicles of infected plants show symptoms. Leaf symptoms begin as chlorosis at the base of the leaf lamina and successively higher leaves show a progression of greater leaf area coverage by the symptoms. Infected chlorotic areas produce massive amounts of asexual spores, generally on the lower surface giving the leaves a 'downy' appearance. Systemically infected plants remain

stunted and either do not produce panicles or produce malformed panicles. In many affected plants 'green ear' symptoms appear on the panicles due to the transformation of floral parts into leafy structures that may be total or partial and such plants do not produce seeds or produce very few seeds. The infected leaves produce sexual spores (oospores) in the necrotic leaf tissue late in the season.

Currently, in India, about 50% of the 9 million ha under pearl millet cultivation is grown with more than 70 hybrids in which DM incidence has been





highly variable, with some hybrids showing more than 90% incidence at farmer's fields. This disease can be assuming alarming levels when a single genetically uniform pearl millet cultivar is repeatedly and extensively grown in a region. Yield losses within the region can reach 30-40%. Needs to be done to establish the role of cuticle thickness and composition on water loss, both before and after stomatal closure. Asynchronous tiller development Plasticity in tiller development is a characteristic of both pearl millet and its wild progenitors (Vadez et al., 2012), with primary tillers developing in the axils of leaves of the main culm and secondary tillers from buds in the axils of leaves on the primary tillers. These tillers transition to flowering at different times, depending on their age. This is called asynchronous tillering and can allow plants to recover from drought by developing tillers that flower

after the dry spell has ended (Craufurd and Bidinger, 1988). A mid-season drought that terminates at or before flowering causes a significant increase in tiller development, which later can fully compensate for grain number and yield losses in the main culm inflorescence (Mahalakshmi et al., 1987). However, drought loss caused during the grain-filling period of pearl millet with terminal drought stress cannot be effectively compensated by this trait (Mahalakshmi et al., 1987), possibly because those tillers initiated and commencing growth under drought stress are unlikely to be productive (Mahalakshmi et al., 1988).

Flowering plasticity Early flowering is an important drought escape mechanism in plants, and in wheat, early flowering after a relatively short vegetative period is a response for impending conditions of terminal stress (Shavrukov et al., 2017). Pearl millet is a short-day plant, but large genotypic differences exist in the photoperiodic requirement for flowering among pearl millet varieties depending upon the zone of latitude that they are adapted to. Almost 54.4% of total cultivated pearl millet germplasms have been found to flower irrespective of the day length, although most are facultatively photoperiod sensitive, showing a delay in flowering time with increasing day length (Rai et al., 1999). Germplasms originating from equatorial Africa are

usually strictly photoperiod sensitive, needing comparatively short days to flower. Landraces with early flowering (33–40 days) are predominantly found in Pakistan, Ghana, Togo, and India; with very late flowering (121–159 days) in Sierra Leone and the Central African Republic (Upadhyaya et al., 2007). Pearl millet grown under long day length together with midseason water stress was found to delay its flowering time but did not cause a significant effect on grain yield (Mahalakshmi et al., 1987). They suggested that later floral initiation is a drought escape mechanism adapted by pearl millet in early mid-season droughts, just as early floral initiation is adaptive for late-season drought. Varieties are adapted to time flowering close to the end of the rainy season, ensuring completion of maturation with remaining soil moisture (Vadez et al., 2012). However, climate change involving changes in rainfall patterns may mean that formerly adapted landraces no longer flower in sync with local conditions. QTL associated with the change in flowering time under drought stress have been found on LG2, colocalized with a major drought tolerant QTL (Tables 1, S1). Co-localized genes included transcription factors belonging to known flowering time gene families, including zinc finger CCH type and MADS-box gene families (Sehgal et al., 2012). 3 Grain yield under drought stress Many genetic analyses

have targeted variation in grain yield in pearl millet, which is a combination of number of productive tillers, size of the inflorescence, fertilization success, and effectiveness of seed development and maturation. 4 Insight into pearl millet drought tolerant mechanisms using transcriptomics the recent availability of the pearl millet draft genome has accelerated the transcriptomic study of drought stress tolerance (Varshney et al., 2017). Five papers have reported transcriptomics study on drought stress in pearl millet (Choudhary and Padaria, 2015; Dudhate et al., 2018; Jaiswal et al., 2018; Shivhare et al., 2020a; Shivhare et al., 2020b). The earliest transcriptomic studies exploring the genes responsible for drought tolerance were done using the suppressive subtraction hybridization technique on seedlings exposed to various stresses, including salt, drying, and cold (Mishra et al., 2007), and 30% polyethylene glycol 600 (Choudhary and Padaria, 2015).

**Conclusions:**

Pearl millet seems to be the solution for a future when temperatures will soar. Decoding and sequencing the pearl millet grain by a team comprising 65 scientists from across 30 research institutions have proved its adaptive capacity and increasing tolerance to drought. This research has been published in the journal, *Nature Biotechnology*. Research coordinated by



the International Crops Research Institute for Semi-Arid Tropics, India, BGI-Shenzhen, China and the French National Research Institute for Sustainable Development used the latest innovations in DNA sequencing and analysis to identify new genetic tools like molecular markers related to drought and heat tolerance, as well as other important traits like better nutrition profile and pest resistance. This will help farmers grow the crop better in terms of productivity, as pearl millet is a staple food crop for millions of people living in the arid and semi-arid areas of Africa and Asia. As experts suggest, both these continents will see increasing incidents of droughts and high temperatures in the coming years due to climate change. Currently, pearl millet is grown on about 27 million hectares worldwide and is a daily food for more than 90 million people. It is also an important source of fodder for millions of farms. However, its yields have remained low over the last six decades, as the cereal is grown on poor soil.

**References:**

1. Alexandratos, N. (2009). World food and agriculture to 2030/50. In Highlights and views from

MID-2009: Paper for the Expert Meeting on Plant Biol.

2. Babiychuk, E., Bouvier-Navé, P., Compagnon, V., Suzuki, M., Muranaka, T., Van Montagu, M., et al. (2008). Allelic mutant series reveal distinct functions for arabidopsis cycloartenol synthase 1 in cell viability and plastid biogenesis. *Proc. Natl. Acad. Sci.*
3. Banerjee, A., and Roychoudhury, A. (2016). Group II late embryogenesis abundant (LEA) proteins: Structural and functional aspects in plant abiotic stress. *Plant Growth Regul.* 79, 1–17., C., Gouzd, Z. Steele, H. P., and Imperio, R. M. (2010). A mutation in GDPmannose pyrophosphorylase causes conditional hypersensitivity to ammonium, resulting in arabidopsis root growth inhibition, altered ammonium metabolism, and hormone homeostasis. *J.*
4. Bidinger, F., Mahalakshmi, V., and Rao, G. D. P. (1987). Assessment of drought resistance in pearl millet (*Pennisetum americanum* (L.) leeke).